

available at www.sciencedirect.comwww.elsevier.com/locate/brainresrev
**BRAIN
RESEARCH
REVIEWS**

Review

Zebrafish and motor control over the last decade

Joseph R. Fetcho^{a,*}, Shin-ichi Higashijima^b, David L. McLean^a

^aW103 Mudd Hall, Department of Neurobiology and Behavior, Cornell University, Ithaca, NY 14853, USA

^bNational Institutes of Natural Sciences, Okazaki Institute for Integrative Bioscience and National Institute for Physiological Sciences, Myodaiji Okazaki, Aichi 444-8787, Japan

ARTICLE INFO

Article history:

Accepted 20 June 2007

Available online 27 July 2007

Keywords:

Zebrafish

Interneuron

Motoneuron

Spinal cord

Transgenic

Mutant

Transcription factor

Recruitment

ABSTRACT

The combination of transparency and accessible genetics is making zebrafish an increasingly important model in studies of motor control. Much of the work on the model has been done over the past decade. Here we review some of the highlights of this work that serve to reveal both the power of the model and its prospects for providing important future insights into the links between neural networks and behavior.

© 2007 Elsevier B.V. All rights reserved.

Contents

1. Introduction	87
2. General aspects of motor behavior.	87
3. Basic biology of spinal networks	87
4. Imaging and ablation studies to reveal activity patterns and functional roles of neurons.	88
5. Mutants and genetic perturbations	89
5.1. Synaptic function	89
5.2. Developmental/pathfinding errors	90
5.3. Spinal injury	90
6. The future	91
Acknowledgments	91
References.	91

* Corresponding author. Fax: +1 607 254 1303.

E-mail address: jrf49@cornell.edu (J.R. Fetcho).

1. Introduction

The zebrafish model is a relative newcomer to studies of motor control and neuronal networks in general. Nonetheless, it has a combination of strengths that are allowing studies that are very difficult or impossible in other models. The two major assets of the zebrafish model are its transparency and its genetic accessibility. The transparency allows optical approaches to the study of structure and function in the intact animal. The genetics allows for the production of both mutant animals with motor disruptions and transgenic fish. This combination of transparency and genetics is unavailable in other vertebrate models (although nematodes offer many of the same strengths in an invertebrate) and is what has driven studies of motor control using zebrafish, almost all of which have occurred in the past decade. Here we review some of the major work, with an eye to further questions in motor control that are made accessible by this model.

2. General aspects of motor behavior

The major motor behaviors and their development have been described in some detail for larval zebrafish (Budick and O'Malley, 2000; Eaton et al., 1977; Saint-Amant and Drapeau, 1998). There is a regular pattern of development with gross body movements followed by the development of rhythmic swimming and, subsequently, slower swimming and pectoral fin movements (Thorsen et al., 2004). Very early movements rely on gap junctional coupling for their production (Saint-Amant and Drapeau, 2001). Later refined movements are used by the larvae during routine swimming, after it has developed a swim bladder to allow it to move from the bottom up into the water column (Borla et al., 2002). Remarkably the larval fish can swim over a very large range of bending frequencies from roughly 15 Hz to 100 Hz (Budick and O'Malley, 2000; Muller and van Leeuwen, 2004). Escape bends are extremely fast with the head bending to touch the tail in about 10 ms, at a rate of about 25° per ms (Liu and Fetcho, 1999).

An early effort to identify the parts of the nervous system responsible for swimming movements indicated that the early spontaneous movements did not require the hindbrain, but that hindbrain was essential for later touch responses and swimming (Saint-Amant and Drapeau, 1998). Recent work indicates that this result was mistaken and that the fish can generate swimming and touch responses without the brain, even at later stages (Downes and Granato, 2006). In this respect, zebrafish are like other vertebrates, in which isolated spinal cord maintains some ability to produce reflex responses and rhythmic motor output.

3. Basic biology of spinal networks

The interpretation of mutants or perturbations often depends upon an understanding of the normal wiring of the networks. Many of the neuronal types in zebrafish spinal cord have been identified and their wiring is the subject of much current work (Bernhardt et al., 1990; Hale et al., 2001; Higashijima et al., 2004b).

Several factors have contributed to reasonably rapid progress in identifying the cells and circuits in zebrafish. *In vivo* calcium imaging has allowed for the rapid identification of neurons active in a particular behavioral context (Brustein et al., 2003b; Cox and Fetcho, 1996; Fetcho and O'Malley, 1995; O'Malley et al., 1996; Ritter et al., 2001). The ability to do patch recording allows for studies of cellular properties and synaptic connectivity that complements imaging work, in which the activity in groups of neurons can be monitored simultaneously (Drapeau et al., 1999; Higashijima et al., 2004a; Kimura et al., 2006; Masino and Fetcho, 2005). Finally, another major advantage is the similarities to networks in *Xenopus* tadpoles, which are already reasonably well understood based upon decades of study (Roberts, 2000; Roberts et al., 1998). This has allowed more directed questions about cell types whose structure and transmitter phenotype made them good candidates for roles that had already been suggested for similar neurons in *Xenopus*. The progress is further accelerated because of the genetics, which have allowed for *in situ* markers of transmitter phenotype to catalog the transmitters that a neuron probably uses (Higashijima et al., 2004b). This can be combined with GFP labeling to define the structure of the neuron as well as its transmitter properties. Finally, transgenic lines with fluorescent subsets of neurons are leading to animals with color coded nervous systems (Higashijima et al., 2000, 2004a; Kimura et al., 2006). This allows for a combination with calcium indicators to look at the pattern of activity in one cell type, or for targeted patch recording directed at neurons of particular phenotypes based upon their fluorescence.

Neuromodulatory systems in zebrafish have also been the subject of considerable attention because of the possibilities of understanding their contributions to behavior and of developing disease models based upon mutations or drugs that disrupt these systems. Dopaminergic, serotonergic, noradrenergic and histamatergic systems in larvae and/or adults have been described morphologically and some receptors and transporters have been cloned (Boehmler et al., 2004; Kaslin and Panula, 2001; Ma, 2003; McLean and Fetcho, 2004a,b; Wang et al., 2006). Pharmacological studies, mutants, and chemical ablations indicate important roles for neuromodulatory systems, but mechanistic studies to reveal the activity patterns of neuromodulatory neurons and their effects on motor circuits are needed (Anichtchik et al., 2004; Brustein et al., 2003a; Rink and Guo, 2004).

One of the most striking aspects of the zebrafish work is that the genetics have finally allowed the determination of the likely relationships between spinal neuron types in fishes (and amphibians) and those in more complex birds and mammals. Developmental studies in chicks and mice revealed a transcription factor code that directs the development of different neuronal types from different dorsoventral regions in spinal cord (Briscoe et al., 2000). These transcription factors can be used as markers of subsets of neurons. The same transcription factor code operates in fish, so the markers can be used to identify what are probably evolutionarily related cell types in fish and mammals.

This has now been done for two neuronal types. The *engrailed-1* positive neurons are ipsilateral, ascending inhibitory neurons that play roles in sensory gating and controlling the duration of motor bursts (Higashijima et al., 2004a; Li et al., 2004) (Fig. 1). The identity of the neurons as *engrailed-1* positive was




Fig. 1 – Imaging structure and function in vivo with genetic tools. (A) Confocal image showing detailed morphology from an intact living fish of an ascending interneuron labeled with GFP under control of the promoter for the *engrailed-1* gene, which is expressed in ascending inhibitory interneurons in zebrafish. The view is a lateral view with the head to the left. **(B)** Two ascending interneurons expressing the genetic calcium indicator cameleon 2.1 under control of the *engrailed-1* promoter. The calcium response of the neuron marked by the asterisk is shown in panel C during a swimming episode that begins at the break in the plot. The neurons are active during swimming, as indicated by the fluorescence ratio increase.

first shown in zebrafish, and was subsequently demonstrated in *Xenopus* for similar ascending neurons. The *engrailed-1* cells in mammals are a heterogeneous population that includes the Renshaw neurons, which provide feedback inhibition of motoneurons (Sapir et al., 2004; Saueressig et al., 1999). The ascending interneurons in fish and frogs have some features akin to Renshaw cells, but they play additional functional roles as well. The evidence suggests that multifunctional neurons like the *engrailed* cells in fish and frogs gave rise to a series of cell types in mammals, each of which has more restricted functional roles. Primitive ascending inhibitory neurons might have given rise to both Renshaw neurons as well as other *engrailed* positive cells in mammals. The role of the *engrailed* cells in frog tadpoles and fishes in shaping motor output to truncate bursts at high frequencies led to the prediction that disrupting this cell type in mammals might slow the frequency of rhythmic movements (Gosgnach et al., 2006; Higashijima et al., 2004a; Li et al., 2004; Sapir et al., 2004; Saueressig et al., 1999). This prediction was recently supported in experiments in which the *engrailed* neurons were selectively inactivated in mice (Gosgnach et al., 2006).

More recently, the neurons in fish that correspond to the *Chx-10* positive cells in mammals were identified as ipsilateral descending excitatory neurons, many of which are rhythmically active during swimming and probably help to drive motor output at higher swimming speeds (Kimura et al., 2006). We might predict that *Chx-10* positive excitatory neurons in mammals play similar roles in driving motor output and that they would be a good place to search for excitatory elements in the mammalian central pattern generator. The ability to link neurons across species makes the study of zebrafish more broadly relevant, as the results from fish are being used to make predictions about the functional roles of neurons in more complex systems, where it is sometimes difficult to predict what role a class of cell might play in the network. The genetics thus serve as a bridge to unite studies in disparate species and elevate our understanding of both the individual species and their evolution.

4. Imaging and ablation studies to reveal activity patterns and functional roles of neurons

Although there were pioneering studies of the development of motor behavior and the startle response of zebrafish many years ago (Eaton et al., 1977), the utility of the animal as a model expanded as a result of a convergence of new methods to image activity *in vivo* with the tractability of screening for mutants. The development of an approach to backfill neurons *in vivo* with calcium indicators offered the possibility to image activity of individual neurons in intact zebrafish (O'Donovan et al., 1993). This allowed the first imaging of neuronal activity with single cell resolution in an intact vertebrate (Fetcho and O'Malley, 1995). The first work was largely technical, both documenting the ability to reliably image groups of motoneurons *in vivo* during an escape response and showing how massive the recruitment of motoneurons is during the escape bend in fish. Every imaged motoneuron was activated when an escape was triggered by a tap on the head of the fish.

The backfilling was then extended to a broader question centered around the functional organization of vertebrate hindbrains (O'Malley et al., 1996). The hindbrains of all vertebrates are segmentally organized, with segmental domains specified by a homeobox code during development. This segmental organization is evident even in adults, where segmentally repeated cell types are evident (Metcalf et al., 1986). The presence of segmentally repeated neurons is thought to result from segmental duplication events during evolution. Their presence raised obvious questions about the functional relationships of the segmental neurons. These neurons could be backfilled with calcium indicators in zebrafish, so their activity in relation to behavior could be studied. Early work examined the activity pattern of a repeated set of reticulospinal neurons thought to be involved in the escape response of fish. This work revealed that the pattern of activity in the serially repeated set of neurons (the Mauthner cell, and MiD2 cm, and MiD3 cm) varied depending on the location of the stimulus used to elicit an

escape. This variation is related to variability in the behavior, with stronger escape bends produced in response to stimuli at the head associated with activation of the entire serial set of cells and the weaker escapes produced by stimuli to the tail involving only the activation of the Mauthner cell. The implication of this work is that serially repeated hindbrain neurons may form functional groups associated with particular behaviors. The pattern of activity in the group may help to determine the different forms of the behavior. There are many serially repeated groups of hindbrain neurons—whether other sets also form functional groups remains to be established.

The contribution of the hindbrain neurons was later tested by focusing a laser beam into the transparent larval fish to kill particular neurons in the hindbrain array and then examine the resulting behavioral deficits (Liu and Fetcho, 1999). The evidence supported the conclusion that the hindbrain neurons contributed differentially to escapes to head and tail stimuli because deleting the Mauthner cell had a dramatic effect on escapes to tail stimuli and none on escapes produced by a head stimulus. This utility of zebrafish for both imaging activity simultaneously in groups of neurons and for testing their contribution to behavior by optical (or genetic, see below) perturbation is perhaps the major advantage of the model. Other models such as flies and mice allow for one or the other, but both are possible throughout the brain and spinal cord of zebrafish.

Imaging studies subsequently revealed a massive activation of hindbrain neurons in response to taps on the head, suggesting a widespread involvement of the reticulospinal system in response to the stimulus (Gahtan et al., 2002). In this and the earlier work on the hindbrain array, the fish was entirely restrained in agar, so the behavior after the stimulus was difficult to discern. One possible explanation of the large reticulospinal activation is that at least some of the neurons are involved in swimming and/or struggling responses following the escape. As one might expect, descending hindbrain neurons do apparently contribute to behaviors other than escape. A carefully performed single cell laser ablation study revealed that a pair of neurons in the nucleus of the medial longitudinal fasciculus with dendrites in the optic tectum and projections to spinal cord contribute to feeding motor responses (Gahtan et al., 2005). Ablation of the neurons reduced prey capture success and appeared to affect the ability of the fish to properly orient to the prey. This ability to link even a few neurons to different motor behaviors offers the promise of understanding the contribution of many cell types.

In order to link neurons to movements, it is important to know which movement is occurring when the neurons are active. This can be difficult with imaging because movement interferes with the image collection, which is why early experiments were done in fish that were entirely embedded in agar. The problem has been resolved to some extent by developing a preparation in which the front of the fish is in agar and the tail is free to move (Ritter et al., 2001). Active neurons can be imaged in the front of the fish while the movements of the tail are filmed to determine which behaviors are being produced in association with the neural activity. This was initially used to study which neurons were active in spinal cord during swimming versus escape responses (Ritter et al., 2001). The work showed that descending commissural neurons (MCoDs), which we now know are excitatory, are active during

slow speed movements and another different descending excitatory interneuron class (CiDs) is active in fast escape responses. This partially moving preparation has allowed studies of the patterns of activation of a class of interneuron during escapes which show that the escape is graded largely by changes in the level of activity in the entire pool of interneurons rather than recruitment of inactive cells (Bhatt et al., 2007).

A recent preliminary report indicates that there is actually a switching of spinal interneuron classes as the swimming speed is increased, with excitatory neurons active at slow speeds being inhibited at higher speeds, when another class of excitatory interneurons becomes active (Masino et al., 2005). These studies suggest that we need to be careful about assuming that the circuits that produce slower movements are the same as the ones that produce fast movements, even when both involve the same overall motor pattern (in the case of swimming, alternating bending patterns that propagate from head to tail).

5. Mutants and genetic perturbations

One of the special strengths of zebrafish is the ability to carry out mutagenesis screens and identify animals with single gene disruptions that affect behavior, including motor behaviors. This was recognized early on in the first large scale mutagenesis study in which mutants were identified with disrupted movements not associated with impaired muscle development (Granato et al., 1996). These fell into many categories, including ones that did not respond to a touch at all (e.g. *macho*, *space cadet*), others that fatigued and so could not sustain swimming (*twitch once*, *shocked*), others that contracted both sides at the same time so they looked like an accordion (*accordion*, *bandoneon*, *diwanka*), and still others that moved wildly or inappropriately (*space cadet*, *techno trousers*).

Many of these have now been studied in more depth both from genetic and functional viewpoints. In a handful of cases, the genes have been identified and there is some understanding of how the gene might contribute to the phenotype. The results can be divided into a few broad categories—those mutants in which synaptic function at the neuromuscular junction or in the CNS is altered leading to a phenotype, and those in which there appear to be axonal pathfinding errors that result in a miswiring of networks.

5.1. Synaptic function

Many of the mutants with motor defects have problems with genes associated with the neuromuscular junction, or with processes in the muscle itself. These include mutations of the acetylcholine receptor, such as *Nic1* and *sofa potato*, which cannot move, as well as the immotile mutant *relaxed*, in which a mutation of the dihydropyridine receptor interferes with the calcium release needed for muscle contraction (Ono et al., 2001; Schredelseker et al., 2005; Sepich et al., 1998; Zhou et al., 2006). These are perhaps less important from a motor control point of view, although they can be used to examine aspects of muscle function and provide a convenient source of immotile fish for physiological experiments.

Some of the more interesting mutations are those that produce fish that can move, but do so in an impaired way. For

example, one of these, called *twice once*, is a mutation in rapsyn, a protein that is involved in localization of acetylcholine receptors at the neuromuscular junction (Ono et al., 2002). These mutants show an inability to swim continuously after a stimulus. Morphological observations with labeled alpha bungarotoxin show that acetylcholine receptors are diffusely distributed rather than localized at the synapses. The mutants also show an abnormally fatigable neuromuscular transmission, consistent with the inability to sustain behavior. This mutation is of some clinical significance because human studies have revealed that the same type of mutation leads to human myasthenic syndrome, also characterized by muscle weakness (Ohno et al., 2002).

The accordion class of mutants is a group showing bilateral muscle contractions. Studies of mutants have revealed that there are many ways to make a fish generate accordion-like movements. One is to disrupt the calcium transporter (SERCA) responsible for removal of the calcium from the cytoplasm after muscle contraction (Gleason et al., 2004; Hirata et al., 2004). When this is not working, as in the mutant named *accordion*, the muscles do not relax properly and the result is that the contraction on one side of the fish overlaps with that on the other, generating the phenotype. The accordion class mutant *Zieharmonika* is a mutation in acetylcholinesterase, which would elevate acetylcholine at the synapses (Downes and Granato, 2004). Early work proposed that central deficits in glycinergic transmission could also produce an accordion phenotype because glycinergic inhibition is known to play a critical role in the alternation of contractions on opposite side of the fish during bending. One of the mutants in the accordion class, called *bandoneon*, indeed turns out to be a mutant of a beta subunit of the glycine receptor which disrupts proper inhibitory transmission and leads to co-contraction of opposite sides of the body (Hirata et al., 2005).

Proper glycinergic transmission is also essential to sustain motor behaviors. The mutant *shocked* shows an inability to sustain movement. The mutation is in the glial, glycine transporter-1, and the disruption appears to elevate glycine levels which may allow for an inappropriate rise in the level of inhibition during movements, thereby truncating the movements (Cui et al., 2005, 2004). The mutation secondarily leads to a persistent electrical coupling of muscle fibers that may reinforce the inability to sustain swimming by inappropriately prolonging muscle contractions (Luna et al., 2004).

5.2. Developmental/pathfinding errors

Some mutants lead to pathfinding errors that result in motor phenotypes. These can be informative about cues used in axonal pathfinding as in the case of the mutants *unplugged* and *diwanka*, both of which involve molecules that play a role in guiding motor axons to their targets (Schneider and Granato, 2006; Zhang et al., 2004). More informative from a behavioral viewpoint are those mutants in which central neurons are disrupted in ways that alter circuit function. These also involve changes in genes important for development that may alter cell number or axonal pathfinding. These are informative when the developmental deficit leads to a wiring change that can offer insight into network function. The mutants *deadly seven* and *space cadet* have both offered circuit level insights.

Deadly seven, a mutant of a *notch* gene, leads to extra Mauthner neurons in the hindbrain (Gray et al., 2001; Liu et al., 2003; van Eeden et al., 1996). Usually there is one on each side of the brain, but there can be as many as 5 or so on each side in mutant fish. Interestingly, this has little impact on the motor output in the escape (Liu et al., 2003). A close look at the morphology of the outputs of individual neurons indicates that the extra neurons divide up the synaptic targets of the neurons normally innervated by a single Mauthner cell. This reveals a good deal of plasticity in the nervous system to accommodate variation in cell number, even when such variations are not seen normally. This makes it easier to imagine how a system such as the escape (startle) built with relatively few neurons in fish could evolve into a mammalian startle circuit with much the same wiring, but many more neurons.

The *space cadet* mutant involves a wiring deficit (Lorent et al., 2001). The phenotype of this mutant is the generation of multiple, inappropriate escape bends, often repeated ones to the same side. Central commissures in the hindbrain of this animal are disrupted, including one that involves the axons of spiral fibers, which play a role in controlling the activation of the Mauthner neuron which initiates the startle. Thus, it may be that an axon pathfinding deficit leads to a behavioral phenotype because of a miswiring of a neuron critical for the response.

Another developmental avenue to explore function is to inject constructs into single cell embryos that lead to structural and possibly behavioral alterations that can be studied functionally. One compelling example of this approach was to inject a Hox gene, which leads to a conversion of a neuron that does not normally form a Mauthner cell into an extra Mauthner neuron (Hale et al., 2004). The result is fish with two Mauthner cells, one in the normal hindbrain segment #4 and the other in hindbrain segment #2. This mimics a reversion to what may be a primitive evolutionary state because hindbrain segments are thought to have been duplicates of one another that later diverged. Lesion experiments show that the extra Mauthner cell can substitute for the normal one when the normal one is removed, indicating again that such switches in segmental identity can be readily accomplished without dramatic functional impairments. Whatever rules guide the development of wiring, they allow extra neurons to easily slot into the networks to contribute to behavior.

5.3. Spinal injury

Zebrafish larvae and adults have also been used to explore the problem of regeneration of axons in spinal cord after injury (Becker et al., 1998, 1997). Unlike mammals, some central axons in fish regenerate well, while others do not. Studies of adults have revealed differences in the upregulation of growth associated molecules that are correlated with regenerative capacity of neurons projecting from the brain to the spinal cord (Becker et al., 1998). The ability to image the regeneration directly in larvae has revealed that cyclic AMP can convert a neuron that has failed to regenerate across a lesion into one whose axon grows to cross the lesion (Bhatt et al., 2004). That growth is associated with both the recovery of activity of neurons postsynaptic to the regenerating axon and with the recovery of behavior. Once again the combination of direct imaging with genetic approaches offers a set of tools that will allow for

exploration of the molecules that promote the regenerative process and that might be clinically relevant in humans.

6. The future

The advantages of zebrafish are likely to propel the model further as an animal in which motor behaviors (and others) can be understood at the cellular level. Progress on several fronts will make it an even better preparation for revealing and testing features of motor organization. Recent work, for example, indicates that interneurons used to move slowly differ from those used to move quickly and that there is a topographic organization of motoneurons and excitatory interneurons in spinal cord, with increasing more dorsal neurons recruited as the frequency of bending movements increases during swimming (McLean et al., 2007). These features of organization were tested by perturbations in which more ventral excitatory interneurons were removed to affect slow swimming without altering faster movements (McLean et al., 2007). Genetically encoded calcium indicators offer the possibility of transgenic lines of fish in which activity in particular subsets of cells can be monitored with single neuron resolution. The first vertebrate transgenic calcium indicator line was generated in zebrafish, but the signals from the older generation of genetic calcium indicators were not especially strong (Higashijima et al., 2003). Increasingly better genetically encoded indicators are being generated that will make imaging easier in subsets of neurons targeted in transgenic lines (Mank et al., 2006; Miyawaki et al., 1997; Nagai et al., 2004; Pologruto et al., 2004; Reiff et al., 2005; Tallini et al., 2006). Powerful genetic tools for perturbations of various sorts have also been developed. Some of the most promising for zebrafish are those that use light activated channels that can activate or inhibit neurons (Banghart et al., 2004; Boyden et al., 2005; Han and Boyden, 2007; Lechner et al., 2002; Li et al., 2005; Szobota et al., 2007; Zhang et al., 2007). One can easily imagine using light to turn off or to activate particular sets of cells to test their contribution to behavior. This can be done on the population as a whole, perhaps even by playing back patterns of spikes in individual neurons using brief flashes of light that elicit single spikes in particular cells. This will move the work more into the realm of testing contributions of neurons to both motor patterns and behavior. Finally, it is possible to image even subcellular synaptic level events in larval fish, such as the movements of GFP tagged calcium dependent protein kinase II, a molecule implicated in synaptic plasticity (Gleason et al., 2003). This can be done noninvasively anywhere in the brain or spinal cord of animals in which activity patterns and behavior can also be examined. This unique combination of features may make zebrafish a favored model for linking molecular, cellular, network and behavioral levels in studies of the neuronal basis of behavior.

Acknowledgments

This work was supported by grants from the National Institutes of Health (NRSA 44728, DLM; NS 26539, JRF) and the Ministry of Education, Science, Technology, Sports and Culture of Japan.

REFERENCES

- Anichtchik, O.V., et al., 2004. Neurochemical and behavioural changes in zebrafish *Danio rerio* after systemic administration of 6-hydroxydopamine and 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *J. Neurochem.* 88, 443–453.
- Banghart, M., et al., 2004. Light-activated ion channels for remote control of neuronal firing. *Nat. Neurosci.* 7, 1381–1386.
- Becker, T., et al., 1997. Axonal regrowth after spinal cord transection in adult zebrafish. *J. Comp. Neurol.* 377, 577–595.
- Becker, T., et al., 1998. Readiness of zebrafish brain neurons to regenerate a spinal axon correlates with differential expression of specific cell recognition molecules. *J. Neurosci.* 18, 5789–5803.
- Bernhardt, R.R., et al., 1990. Identification of spinal neurons in the embryonic and larval zebrafish. *J. Comp. Neurol.* 302, 603–616.
- Bhatt, D.H., et al., 2004. Cyclic AMP-induced repair of zebrafish spinal circuits. *Science* 305, 254–258.
- Bhatt, D.H., et al., 2007. Grading movement strength by changes in firing intensity versus recruitment of spinal interneurons. *Neuron* 53, 91–102.
- Boehmler, W., et al., 2004. Evolution and expression of D2 and D3 dopamine receptor genes in zebrafish. *Dev. Dyn.* 230, 481–493.
- Borla, M.A., et al., 2002. Prey capture by larval zebrafish: evidence for fine axial motor control. *Brain Behav. Evol.* 60, 207–229.
- Boyden, E.S., et al., 2005. Millisecond-timescale, genetically targeted optical control of neural activity. *Nat. Neurosci.* 8, 1263–1268.
- Briscoe, J., et al., 2000. A homeodomain protein code specifies progenitor cell identity and neuronal fate in the ventral neural tube. *Cell* 101, 435–445.
- Brusteine, E., et al., 2003a. Serotonin patterns locomotor network activity in the developing zebrafish by modulating quiescent periods. *J. Neurobiol.* 57, 303–322.
- Brusteine, E., et al., 2003b. “In vivo” monitoring of neuronal network activity in zebrafish by two-photon Ca(2+) imaging. *Pflugers Arch.* 446, 766–773.
- Budick, S.A., O’Malley, D.M., 2000. Locomotor repertoire of the larval zebrafish: swimming, turning and prey capture. *J. Exp. Biol.* 203, 2565–2579.
- Cox, K.J.A., Fetcho, J.R., 1996. Labeling blastomeres with a calcium indicator: a non-invasive method of visualizing neuronal activity in zebrafish. *J. Neurosci. Methods* 68, 185–191.
- Cui, W.W., et al., 2004. Shocked Gene is required for the function of a premotor network in the zebrafish CNS. *J. Neurophysiol.* 92, 2898–2908.
- Cui, W.W., et al., 2005. The zebrafish shocked gene encodes a glycine transporter and is essential for the function of early neural circuits in the CNS. *J. Neurosci.* 25, 6610–6620.
- Downes, G.B., Granato, M., 2004. Acetylcholinesterase function is dispensable for sensory neurite growth but is critical for neuromuscular synapse stability. *Dev. Biol.* 270, 232–245.
- Downes, G.B., Granato, M., 2006. Supraspinal input is dispensable to generate glycine-mediated locomotive behaviors in the zebrafish embryo. *J. Neurobiol.* 66, 437–451.
- Drapeau, P., et al., 1999. In vivo recording from identifiable neurons of the locomotor network in the developing zebrafish. *J. Neurosci. Methods* 88, 1–13.
- Eaton, R.C., et al., 1977. Functional development in the Mauthner cell system of embryos and larvae of the zebra fish. *J. Neurobiol.* 8, 151–172.
- Fetcho, J.R., O’Malley, D.M., 1995. Visualization of active neural circuitry in the spinal-cord of intact zebrafish. *J. Neurophysiol.* 73, 399–406.
- Gahtan, E., et al., 2002. Evidence for a widespread brain stem escape network in larval zebrafish. *J. Neurophysiol.* 87, 608–614.

- Gahtan, E., et al., 2005. Visual prey capture in larval zebrafish is controlled by identified reticulospinal neurons downstream of the tectum. *J. Neurosci.* 25, 9294–9303.
- Gleason, M.R., et al., 2003. Translocation of CaM kinase II to synaptic sites in vivo. *Nat. Neurosci.* 6, 217–218.
- Gleason, M.R., et al., 2004. A mutation in *serca* underlies motility dysfunction in accordion zebrafish. *Dev. Biol.* 276, 441–451.
- Gosgnach, S., et al., 2006. V1 spinal neurons regulate the speed of vertebrate locomotor outputs. *Nature* 440, 215–219.
- Granato, M., et al., 1996. Genes controlling and mediating locomotion behavior in the zebrafish embryo and larva. *Development* 123, 399–413.
- Gray, M., et al., 2001. Zebrafish deadly seven functions in neurogenesis. *Dev. Biol.* 237, 306–323.
- Hale, M.E., et al., 2001. A confocal study of spinal interneurons in living larval zebrafish. *J. Comp. Neurol.* 437, 1–16.
- Hale, M.E., et al., 2004. Hox gene misexpression and cell-specific lesions reveal functionality of homeotically transformed neurons. *J. Neurosci.* 24, 3070–3076.
- Han, X., Boyden, E.S., 2007. Multiple-Color Optical Activation, Silencing, and Desynchronization of Neural Activity, with Single-Spike Temporal Resolution. *PLOS one* 2, e299. doi:10.1371/journal.pone.0000299.
- Higashijima, S., et al., 2000. Visualization of cranial motor neurons in live transgenic zebrafish expressing green fluorescent protein under the control of the *islet-1* promoter/enhancer. *J. Neurosci.* 20, 206–218.
- Higashijima, S.I., et al., 2003. Imaging neuronal activity during zebrafish behavior with a genetically encoded calcium indicator. *J. Neurophysiol.* 90, 3986–3997.
- Higashijima, S., et al., 2004a. Engrailed-1 expression marks a primitive class of inhibitory spinal interneuron. *J. Neurosci.* 24, 5827–5839.
- Higashijima, S., et al., 2004b. Neurotransmitter properties of spinal interneurons in embryonic and larval zebrafish. *J. Comp. Neurol.* 480, 19–37.
- Hirata, H., et al., 2004. Accordion, a zebrafish behavioral mutant, has a muscle relaxation defect due to a mutation in the ATPase Ca²⁺ pump *SERCA1*. *Development* 131, 5457–5468.
- Hirata, H., et al., 2005. Zebrafish bandoneon mutants display behavioral defects due to a mutation in the glycine receptor beta-subunit. *Proc. Natl. Acad. Sci. U. S. A.* 102, 8345–8350.
- Kaslin, J., Panula, P., 2001. Comparative anatomy of the histaminergic and other aminergic systems in zebrafish (*Danio rerio*). *J. Comp. Neurol.* 440, 342–377.
- Kimura, Y., et al., 2006. *alx*, a zebrafish homolog of Chx10, marks ipsilateral descending excitatory interneurons that participate in the regulation of spinal locomotor circuits. *J. Neurosci.* 26, 5684–5697.
- Lechner, H.A., et al., 2002. A genetic method for selective and quickly reversible silencing of mammalian neurons. *J. Neurosci.* 22, 5287–5290.
- Li, W.C., et al., 2004. Primitive roles for inhibitory interneurons in developing frog spinal cord. *J. Neurosci.* 24, 5840–5848.
- Li, X., et al., 2005. Fast noninvasive activation and inhibition of neural and network activity by vertebrate rhodopsin and green algae channel rhodopsin. *Proc. Natl. Acad. Sci. U. S. A.* 102, 17816–17821.
- Liu, K.S., Fetcho, J.R., 1999. Laser ablations reveal functional relationships of segmental hindbrain neurons in zebrafish. *Neuron* 23, 325–335.
- Liu, K.S., et al., 2003. Mutations in *deadly seven/notch1a* reveal developmental plasticity in the escape response circuit. *J. Neurosci.* 23, 8159–8166.
- Lorent, K., et al., 2001. The zebrafish space cadet gene controls axonal pathfinding of neurons that modulate fast turning movements. *Development* 128, 2131–2142.
- Luna, V.M., et al., 2004. Persistent electrical coupling and locomotory dysfunction in the zebrafish mutant *shocked*. *J. Neurophysiol.* 92, 2003–2009.
- Ma, P.M., 2003. Catecholaminergic systems in the zebrafish: IV. Organization and projection pattern of dopaminergic neurons in the diencephalon. *J. Comp. Neurol.* 460, 13–37.
- Mank, M., et al., 2006. A FRET-based calcium biosensor with fast signal kinetics and high fluorescence change. *Biophys. J.* 90, 1790–1796.
- Masino, M.A., Fetcho, J.R., 2005. Fictive swimming motor patterns in wild type and mutant larval zebrafish. *J. Neurophysiol.* 93, 3177–3188.
- Masino, M.A., et al., 2005. Identification of an intersegmental interneuron that may drive slow swimming movements in larval zebrafish. *Soc. Neurosci. Abst.* 31.
- McLean, D.L., Fetcho, J.R., 2004a. Ontogeny and innervation patterns of dopaminergic, noradrenergic, and serotonergic neurons in larval zebrafish. *J. Comp. Neurol.* 480, 38–56.
- McLean, D.L., Fetcho, J.R., 2004b. Relationship of tyrosine hydroxylase and serotonin immunoreactivity to sensorimotor circuitry in larval zebrafish. *J. Comp. Neurol.* 480, 57–71.
- McLean, D.L., et al., 2007. A topographic map of recruitment in spinal cord. *Nature* 446, 71–75.
- Metcalfe, W.K., et al., 1986. Segmental homologies among reticulospinal neurons in the hindbrain of the zebrafish larva. *J. Comp. Neurol.* 251, 147–159.
- Miyawaki, A., et al., 1997. Fluorescent indicators for Ca²⁺ based on green fluorescent proteins and calmodulin [see comments]. *Nature* 388, 882–887.
- Muller, U.K., van Leeuwen, J.L., 2004. Swimming of larval zebrafish: ontogeny of body waves and implications for locomotory development. *J. Exp. Biol.* 207, 853–868.
- Nagai, T., et al., 2004. Expanded dynamic range of fluorescent indicators for Ca(2+) by circularly permuted yellow fluorescent proteins. *Proc. Natl. Acad. Sci. U. S. A.* 101, 10554–10559.
- O'Donovan, M.J., et al., 1993. Real-time imaging of neurons retrogradely and anterogradely labelled with calcium-sensitive dyes. *J. Neurosci. Methods* 46, 91–106.
- O'Malley, D.M., et al., 1996. Imaging the functional organization of zebrafish hindbrain segments during escape behaviors. *Neuron* 17, 1145–1155.
- Ohno, K., et al., 2002. Rapsyn mutations in humans cause endplate acetylcholine-receptor deficiency and myasthenic syndrome. *Am. J. Hum. Genet.* 70, 875–885.
- Ono, F., et al., 2001. Paralytic zebrafish lacking acetylcholine receptors fail to localize rapsyn clusters to the synapse. *J. Neurosci.* 21, 5439–5448.
- Ono, F., et al., 2002. The Zebrafish motility mutant *twitch* once reveals new roles for rapsyn in synaptic function. *J. Neurosci.* 22, 6491–6498.
- Pologruto, T.A., et al., 2004. Monitoring neural activity and [Ca²⁺] with genetically encoded Ca²⁺ indicators. *J. Neurosci.* 24, 9572–9579.
- Reiff, D.F., et al., 2005. In vivo performance of genetically encoded indicators of neural activity in flies. *J. Neurosci.* 25, 4766–4778.
- Rink, E., Guo, S., 2004. The too few mutant selectively affects subgroups of monoaminergic neurons in the zebrafish forebrain. *Neuroscience* 127, 147–154.
- Ritter, D.A., et al., 2001. In vivo imaging of zebrafish reveals differences in the spinal networks for escape and swimming movements. *J. Neurosci.* 21, 8956–8965.
- Roberts, A., 2000. Early functional organization of spinal neurons in developing lower vertebrates. *Brain Res. Bull.* 53, 585–593.
- Roberts, A., et al., 1998. Central circuits controlling locomotion in young frog tadpoles. *Ann. N.Y. Acad. Sci.* 860, 19–34.
- Saint-Amant, L., Drapeau, P., 1998. Time course of the development of motor behaviors in the zebrafish embryo. *J. Neurobiol.* 37, 622–632.
- Saint-Amant, L., Drapeau, P., 2001. Synchronization of an embryonic network of identified spinal interneurons solely by electrical coupling. *Neuron* 31, 1035–1046.
- Sapir, T., et al., 2004. Pax6 and engrailed 1 regulate two distinct aspects of renshaw cell development. *J. Neurosci.* 24, 1255–1264.

- Saueressig, H., et al., 1999. Engrailed-1 and netrin-1 regulate axon pathfinding by association interneurons that project to motor neurons. *Development* 126, 4201–4212.
- Schneider, V.A., Granato, M., 2006. The myotomal diwanka (lh3) glycosyltransferase and type XVIII collagen are critical for motor growth cone migration. *Neuron* 50, 683–695.
- Schredelseker, J., et al., 2005. The beta 1a subunit is essential for the assembly of dihydropyridine-receptor arrays in skeletal muscle. *Proc. Natl. Acad. Sci. U. S. A.* 102, 17219–17224.
- Sepich, D.S., et al., 1998. An altered intron inhibits synthesis of the acetylcholine receptor alpha-subunit in the paralyzed zebrafish mutant *nic1*. *Genetics* 148, 361–372.
- Szobota, S., 2007. Remote control of neuronal activity with a light-gated glutamate receptor. *Neuron* 54, 535–545.
- Tallini, Y.N., et al., 2006. Imaging cellular signals in the heart in vivo: cardiac expression of the high-signal Ca²⁺ indicator GCaMP2. *Proc. Natl. Acad. Sci. U. S. A.* 103, 4753–4758.
- Thorsen, D.H., et al., 2004. Swimming of larval zebrafish: fin-axis coordination and implications for function and neural control. *J. Exp. Biol.* 207, 4175–4183.
- van Eeden, F.J., et al., 1996. Mutations affecting somite formation and patterning in the zebrafish, *Danio rerio*. *Development* 123, 153–164.
- Wang, Y., et al., 2006. Characterization and expression of serotonin transporter genes in zebrafish. *Tohoku J. Exp. Med.* 208, 267–274.
- Zhang, J., et al., 2004. Zebrafish unplugged reveals a role for muscle-specific kinase homologs in axonal pathway choice. *Nat. Neurosci.* 7, 1303–1309.
- Zhang, F., et al., 2007. Multimodal fast optical interrogation of neural circuitry. *Nature* 446, 633–639.
- Zhou, W., et al., 2006. Non-sense mutations in the dihydropyridine receptor beta1 gene. *CACNB1*, paralyze zebrafish relaxed mutants. *Cell Calcium* 39, 227–236.